

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204 Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 30 October 1996		3. REPORT TYPE AND DATES COVERED Technical
4. TITLE AND SUBTITLE SYNTHESIS AND CHARACTERIZATION OF AN ORGANOTHALLIUM-PHOSPHORUS ADDUCT: CRYSTAL STRUCTURE OF $(\text{Me}_3\text{SiCH}_2)_3\text{TI}\cdot\text{P}(\text{SiMe}_3)_3$			5. FUNDING NUMBERS •N00014-95-1-0194 R&T Project 3135008---16 •Dr. Harold E. Guard	
6. AUTHOR(S) Ryan A. Baldwin, Richard L. Wells, and Peter S. White				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry Duke University Durham, NC 27708-0346			8. PERFORMING ORGANIZATION REPORT NUMBER Technical Report No. DU/DC/TR-59	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 300 North Quincy Street Arlington, VA 22217-5000			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Accepted for publication in <i>Main Group Chemistry</i>			19961209 040	
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The organothallium phosphorus adduct $(\text{Me}_3\text{SiCH}_2)_3\text{TI}\cdot\text{P}(\text{SiMe}_3)_3$ (1) was prepared by combining $(\text{Me}_3\text{SiCH}_2)_3\text{TI}$ and $\text{P}(\text{SiMe}_3)_3$ at room temperature. Compound 1 was characterized by ^1H , $^{13}\text{C}\{^1\text{H}\}$, and $^{31}\text{P}\{^1\text{H}\}$ NMR, partial elemental analysis, EI mass spectrometry, and single-crystal X-ray analysis, the first to be reported for a thallium-group 15 adduct. Crystal data for 1 : trigonal system, space group $P\bar{3}_1$, with $a = 16.063(6)$ Å, $c = 12.148(3)$ Å, $D_{\text{calcd}} = 1.315$ g cm $^{-3}$, and $V = 2714.3(11)$ Å 3 for $Z = 3$. Refinement converged at $R = 0.042$ ($R_w = 0.045$). The TI-P bond length in 1 , previously unreported for compounds of this type, was found to be $2.922(3)$ Å.				
14. SUBJECT TERMS thallium, phosphorus, adduct, synthesis, structure			15. NUMBER OF PAGES 14	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

OFFICE OF NAVAL RESEARCH

Grant N00014-95-1-0194
R&T Project 3135008---16

Dr. Harold E. Guard

Technical Report No. DU/DC/TR-59

**SYNTHESIS AND CHARACTERIZATION OF AN
ORGANOTHALLIUM-PHOSPHORUS ADDUCT:
CRYSTAL STRUCTURE OF $(\text{Me}_3\text{SiCH}_2)_3\text{Ti}\cdot\text{P}(\text{SiMe}_3)_3$**

RYAN A. BALDWIN,¹ RICHARD L. WELLS,¹ AND PETER WHITE²

¹Department of Chemistry, Duke University, Durham, NC 27708

²Department of Chemistry, University of North Carolina, Chapel Hill, NC 27514

Accepted for publication in *Main Group Chemistry*

Duke University
Department of Chemistry,
P. M. Gross Chemical Laboratory
Box 90346
Durham, NC 27708-0346

30 October 1996

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

**SYNTHESIS AND CHARACTERIZATION OF AN ORGANOThALLIUM-
PHOSPHORUS ADDUCT: CRYSTAL STRUCTURE OF $(\text{Me}_3\text{SiCH}_2)_3\text{Tl}\cdot\text{P}(\text{SiMe}_3)_3$**

RYAN A. BALDWIN AND RICHARD L. WELLS*

Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke University,
Durham, North Carolina 27708, U S A.

PETER S. WHITE

Department of Chemistry, Venable Hall, University of North Carolina at Chapel Hill,
Chapel Hill, North Carolina 27514, U S A.

(Revised August 20, 1996; accepted August 8, 1996)

Abstract

The organothallium phosphorus adduct $(\text{Me}_3\text{SiCH}_2)_3\text{Tl}\cdot\text{P}(\text{SiMe}_3)_3$ (**1**) was prepared by combining $(\text{Me}_3\text{SiCH}_2)_3\text{Tl}$ and $\text{P}(\text{SiMe}_3)_3$ at room temperature. Compound **1** was characterized by ^1H , $^{13}\text{C}\{^1\text{H}\}$, and $^{31}\text{P}\{^1\text{H}\}$ NMR, partial elemental analysis, EI mass spectrometry, and single-crystal X-ray analysis, the first to be reported for a thallium-group 15 adduct. Crystal data for **1**: trigonal system, space group $P\bar{3}_1$, with $a = 16.063(6)$ Å, $c = 12.148(3)$ Å, $D_{\text{calcd}} = 1.315$ g cm $^{-3}$, and $V = 2714.3(11)$ Å 3 for $Z = 3$. Refinement converged at $R = 0.042$ ($R_w = 0.045$). The Tl-P bond length in **1**, previously unreported for compounds of this type, was found to be 2.922(3) Å.

INTRODUCTION

As a result of our efforts to find facile methods of forming 13-15 bonds, we have prepared and characterized a variety of new and interesting compounds¹. Among these are Lewis acid-base

* Author for correspondence

adducts of the type $R_3M \cdot E(SiMe_3)_3$ ($R = Me_3SiCH_2$, $M = Ga$, $E = As$ or P ; ^{1c} $M = In$, $E = As$ ^{1a} or P ^{1c}; $R = Ph$ ^{1a,b}, $M = Ga$, $E = As$ or P). The successful isolation of these compounds motivated us to continue our studies of 13-15 adducts, specifically those containing thallium. Although thallium-group 15 adducts have been prepared², no solid-state structures of such compounds have been reported.

To this end, herein, we report the synthesis and structural characterization of the novel organothallium phosphorus compound $(Me_3SiCH_2)_3Tl \cdot P(SiMe_3)_3$ (**1**).

EXPERIMENTAL

Synthesis

General Considerations. All manipulations of air- and moisture-sensitive materials were performed in a Vacuum Atmospheres HE-493 Dri-Lab containing an argon atmosphere or by general Schlenk techniques. Pentane and hexane were distilled over sodium-potassium alloy under dry dinitrogen. $(Me_3SiCH_2)_3Tl$ ³, and $P(SiMe_3)_3$ ⁴ were synthesized by literature procedures. ¹H, ¹³C{¹H}, and ³¹P{¹H} NMR spectra were recorded on a Varian Unity 400 spectrometer operating at 400, 100.6, and 161.9 MHz, respectively. ¹H and ¹³C{¹H} spectra were referenced to TMS by using the residual protons or carbons of deuterated benzene at δ 7.15 or 128 ppm, respectively. ³¹P{¹H} spectra were referenced externally to 85% H₃PO₄ at δ 0.00 ppm. All NMR samples were prepared in 5-mm tubes which were septum-sealed under argon. Melting points (uncorrected) were obtained with a Thomas-Hoover Uni-melt apparatus and capillaries were flame-sealed under argon. Elemental Analyses were performed by E+R Microanalytical Laboratory, Inc., Corona, NY. Mass spectral data were collected on a JEOL JMS-SX 102A spectrometer operating in the electron ionization mode at 20 eV. X-ray crystallographic data were obtained at -120 °C on a Rigaku AFC6/S diffractometer utilizing graphite-monochromated Mo-K α ($\lambda = 0.71073$ Å) radiation.

Preparation of $(\text{Me}_3\text{SiCH}_2)_3\text{Tl}\cdot\text{P}(\text{SiMe}_3)_3$ (**1**)

$(\text{Me}_3\text{SiCH}_2)_3\text{Tl}$ (0.466 g, 1.00 mmol) dissolved in 25 mL of pentane was added to a 250 mL round-bottomed screw-top flask equipped with a Teflon valve and magnetic stirbar. $\text{P}(\text{SiMe}_3)_3$ (0.251 g, 1.00 mmol) dissolved in 25 mL of pentane was added dropwise to the stirred $(\text{Me}_3\text{SiCH}_2)_3\text{Tl}$ solution. The resulting clear solution was stirred at room temperature outside of the dry box for 24 h. The volatiles were removed *in vacuo* to yield a faint-purple crystalline solid which was extracted into 5 mL of warm hexane. Cooling of the extract to $-30\text{ }^\circ\text{C}$ for 2 d afforded colorless X-ray quality crystals of **1** (0.562 g, 78.4%). mp. $68\text{--}71\text{ }^\circ\text{C}$. Anal. Calcd. (found) for $\text{C}_{21}\text{H}_{60}\text{PSi}_6\text{Tl}$: C, 35.20 (35.09); H, 8.44 (8.20). ^1H NMR (C_6D_6): δ 0.08, 0.90 [d, $-\text{CH}_2$, $^2J_{\text{Tl-H}} = 329\text{ Hz}$], 0.13 [d, $-\text{SiMe}_3$, $^4J_{\text{Tl-H}} = 7\text{ Hz}$], 0.33 [d, P-SiMe_3 , $^3J_{\text{P-H}} = 5\text{ Hz}$]. $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ 2.07, 3.01 [d, $-\text{CH}_2\text{Si}(\underline{\text{CH}}_3)_3$, $^3J_{\text{Tl-C}} = 95\text{ Hz}$], 4.13 [d, $-\text{PSi}(\underline{\text{CH}}_3)_3$, $^2J_{\text{P-C}} = 12\text{ Hz}$], 29.02, 37.32 [d, $-\underline{\text{CH}}_2\text{Si}(\text{CH}_3)_3$, $^1J_{\text{Tl-C}} = 835\text{ Hz}$]. $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ -251.7 (s). Mass spec.: m/z 451, $(\text{C}_{11}\text{H}_{33}\text{Si}_3\text{Tl})^+$; 379, $(\text{C}_8\text{H}_{22}\text{Si}_2\text{Tl})^+$; 250, $(\text{C}_9\text{H}_{27}\text{PSi})^+$.

X-Ray Crystal Structure Determination

Crystallographic data for **1** are summarized in Table 1. The crystal used was a colorless block which was mounted on a glass fiber with a viscous oil under a stream of cold dinitrogen. X-ray intensity data were recorded at $-120\text{ }^\circ\text{C}$ and the structures were solved by direct methods. Full-matrix least-squares refinement with weights based upon counter statistics was performed. Hydrogen atoms were incorporated at their calculated positions using a riding model in the later iterations of refinement which converged at $R = 0.042$ ($R_w = 0.045$). A final difference-Fourier synthesis revealed no unusual features (max. 1.64 , min. $-1.24\text{ e } \text{\AA}^{-3}$). Crystallographic calculations were performed using the NRCVAX⁵ suite of structure determination programs. For all structure-factor calculations, neutral atom scattering factors and their anomalous dispersion corrections were taken from ref. 6. Interatomic distances and angles are given in Table 2. Non-hydrogen atom fractional atomic coordinates are listed in Table 3. An ORTEP⁷

diagram showing the solid-state conformation and atom numbering scheme of **1** is presented in the Figure.

RESULTS AND DISCUSSION

The 1:1 mole ratio reaction of $(\text{Me}_3\text{SiCH}_2)_3\text{Tl}$ with $\text{P}(\text{SiMe}_3)_3$ at room temperature affords the adduct $(\text{Me}_3\text{SiCH}_2)_3\text{Tl}\cdot\text{P}(\text{SiMe}_3)_3$ (**1**) in a good yield. Compound **1** is a crystalline material which is stable over long periods of time under inert atmosphere. In addition, **1** is surprisingly stable upon exposure to air and moisture and does not decompose rapidly. Interestingly, the resonances observed in the ^1H , $^{13}\text{C}\{^1\text{H}\}$, and $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of **1** have chemical shifts that are very close to those observed for the starting materials, which suggests that **1** dissociates in benzene solution at room temperature. The ^1H and $^{13}\text{C}\{^1\text{H}\}$ spectra also evidence the fact that the thallium atom, with its spin of 1/2, couples to the protons and carbon atoms of its three Me_3SiCH_2 - ligands (see references 2a and 2b for representative ^1H and ^{13}C spectral parameters of organothallium compounds).

Compound **1** crystallizes in the trigonal system with three molecules occupying the general positions of the $\text{P}3_1$ space group. In the solid state, this molecule adopts the conformation depicted in the Figure. The most noteworthy structural feature of **1** is the novel Tl-P bond length of 2.922(3) Å. This bond is longer than the sum of the covalent radii⁸ (2.62 Å) for the thallium and phosphorus atoms, which is not surprising due to the dative nature of the bond. The coordination geometry about the thallium center is that of a distorted tetrahedron with a mean C-Tl-C angle of 118.0° and a mean C-Tl-P angle of 98.1°. Mean bond angles at the P atom [$\text{Tl-P-Si} = 112.5^\circ > \text{Si-P-Si} = 106.3^\circ$] reflect the relayed effect of the steric compressions resulting from the angular deformations around Tl. Since Tl-P analogs of **1** are not known, comparison of its bond lengths and bond angles with similar compounds is confined to the analogous gallium and indium compounds $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}\cdot\text{P}(\text{SiMe}_3)_3$ (**2**)^{1c} and $(\text{Me}_3\text{SiCH}_2)_3\text{In}\cdot\text{P}(\text{SiMe}_3)_3$ (**3**)^{1c}. As expected, the metal-phosphorus bond lengths of 2.646(3) and 2.771(2) Å in **2** and **3**, respectively, are significantly shorter than the observed thallium-

phosphorus linkage in **1**. Surprisingly, the observed degree of angular distortion around the thallium center in **1** is in accord with that seen in **2** [mean C-Ga-C angle = 116.7°, mean C-Ga-P angle = 100.3°] and **3** [mean C-In-C angle = 116.2°, mean C-In-P angle = 101.4°] despite the metal size difference.

Future investigations in this area will focus on determining the utility of **1** as a single-source precursor to TlP and on preparing alternate organothallium pnictogen compounds.

ACKNOWLEDGEMENTS

We wish to thank the Office of Naval Research for their generous financial support. R. A. B. wishes to thank the Lucent Technologies Bell Laboratories Cooperative Research Fellowship Program and the Duke Endowment Fellowship Program for support.

SUPPLEMENTARY DATA


Additional material consisting of a summary of crystallographic data and measurements, atomic coordinates, thermal parameters, bond distances and angles, and structure factors (9 pages).

REFERENCES

1. See for example the following: (a) Wells, R.L.; McPhail, A. T.; Jones, L. J., III; Self, M. F.; Butcher, R. J. *Organometallics* **1992**, 11, 2694. (b) Wells, R. L.; Self, M. F.; Aubuchon, S. R.; Woudenberg, R. C.; Jasinski, J. P.; Butcher, R. J. *Organometallics* **1992**, 11, 3370. (c) Wells, R. L.; Baldwin, R. A.; White, P.S. *Organometallics* **1995**, 14, 2123. (d) Wells, R. L.; Aubuchon, S. R.; Kher, S. S.; Lube, M. S.; White, P.S. *Chem. Mater.* **1995**, 7, 793.
2. (a) Coates, G. E.; Whitcombe, R. A. *J. Chem. Soc.* **1956**, 3351. (b) Maher, J. P.; Evans, D. F. *J. Chem. Soc.* **1963**, 5534.
3. Brady, F.; Henrick, K.; Matthews, R. W.; Gillies, D. G. *J. Organomet. Chem.* **1980**, 193, 21.
4. Becker, G.; and Hoelderich, W. *Chem. Ber.* **1975**, 108, 2484.
5. Gabe, E. J.; Page, Y. L.; Charland, J. P.; Lee, F. L.; and White, P. S. *J. Appl. Cryst.* **1989**, 22, 384.
6. *International Tables for X-ray Crystallography*; The Kynoch Press, Birmingham, England, Vol. IV, 1974.
7. Johnson, C. K. ORTEP- A Fortran Thermal Ellipsoid Plot Program, Technical Report ORNL-5138, Oak Ridge, 1976.
8. Emsley, J. in *The Elements*, 2nd edn., Clarendon Press, Oxford, 1991.

Caption to Figure

Figure. ORTEP diagram (30% probability ellipsoids) showing the solid state structure and atom numbering scheme of **1**. Hydrogen atoms are omitted for clarity.



Figure

Table 1. Crystallographic Data and Measurements for $(\text{Me}_3\text{SiCH}_2)_3\text{Ti}\cdot\text{P}(\text{SiMe}_3)_3$ (**1**)

	1
molecular formula	$\text{C}_{21}\text{H}_{60}\text{PSi}_6\text{Ti}$
formula weight	716.55
crystal system	trigonal
space group	$P3_1$
a, Å	16.063(6)
c, Å	12.148(3)
V, Å ³	2714.3(11)
Z	3
radiation (wavelength, Å)	Mo-K α (0.71073)
μ , mm ⁻¹	4.75
temp, °C	-120
D _{calcd} , g cm ⁻³	1.315
crystal dimens., mm	0.30 x 0.30 x 0.30
T _{max} ; T _{min}	0.360:0.300
scan type	ω
2 Θ _{max} , deg	50
no. of rflns recorded	3998
no. of non-equiv.	3998
rflns recorded	
no. of rflns retained,	3333
I > 2.5 σ (I)	

Table 1 (continued)

	1
no. of params.	262
refined	
R; R _w ^a	0.042; 0.045
goodness-of-fit ^b	1.21
max shift / esd. in final	0.012
least-squares cycle	
final max, min $\Delta\rho$, e/Å ⁻³	1.64; -1.24

$$^a R = \Sigma(|F_o| - |F_c|) / \Sigma |F_o| ; R_w = [\Sigma w_i (|F_o| - |F_c|)^2 / \Sigma w_i |F_o|^2]^{1/2}.$$

$$^b \text{Goodness-of-fit} = [\Sigma w_i \Delta^2 / (N_{\text{observations}} - N_{\text{parameters}})]^{1/2}.$$

Table 2. Bond Distances (Å) and bond angles (°) for **1**, with Estimated Standard Deviations in Parentheses

Bond Lengths			
Tl(1)-P(1)	2.922(3)	Si(3)-C(32)	1.87(2)
Tl(1)-C(41)	2.24(1)	Si(3)-C(33)	1.84(2)
Tl(1)-C(51)	2.23(2)	Si(4)-C(41)	1.85(2)
Tl(1)-C(61)	2.25(2)	Si(4)-C(42)	1.85(2)
P(1)-Si(1)	2.267(5)	Si(4)-C(43)	1.87(2)
P(1)-Si(2)	2.249(5)	Si(4)-C(44)	1.85(2)
P(1)-Si(3)	2.261(5)	Si(5)-C(51)	1.86(2)
Si(1)-C(11)	1.85(2)	Si(5)-C(52)	1.88(2)
Si(1)-C(12)	1.86(2)	Si(5)-C(53)	1.87(2)
Si(1)-C(13)	1.84(2)	Si(5)-C(54)	1.86(2)
Si(2)-C(21)	1.84(2)	Si(6)-C(61)	1.82(1)
Si(2)-C(22)	1.86(2)	Si(6)-C(62)	1.87(2)
Si(2)-C(23)	1.84(2)	Si(6)-C(63)	1.86(2)
Si(3)-C(31)	1.87(2)	Si(6)-C(64)	1.85(2)
Bond Angles			
P(1)-Tl(1)-C(41)	98.9(4)	P(1)-Si(3)-C(33)	107.7(5)
P(1)-Tl(1)-C(51)	95.1(4)	C(31)-Si(3)-C(32)	108.9(7)
P(1)-Tl(1)-C(61)	100.4(4)	C(31)-Si(3)-C(33)	109.4(7)
C(41)-Tl(1)-C(51)	116.5(5)	C(32)-Si(3)-C(33)	109.1(7)
C(41)-Tl(1)-C(61)	116.2(5)	C(41)-Si(4)-C(42)	109.3(7)
C(51)-Tl(1)-C(61)	121.1(5)	C(41)-Si(4)-C(43)	110.6(7)
Tl(1)-P(1)-Si(1)	110.7(2)	C(41)-Si(4)-C(44)	111.5(7)
Tl(1)-P(1)-Si(2)	113.7(2)	C(42)-Si(4)-C(43)	109.0(8)
Tl(1)-P(1)-Si(3)	113.1(2)	C(42)-Si(4)-C(44)	108.5(8)
Si(1)-P(1)-C(11)	105.8(2)	C(43)-Si(4)-C(44)	107.9(9)
Si(1)-P(1)-C(12)	107.1(2)	C(51)-Si(5)-C(52)	108.5(7)
Si(2)-P(1)-C(13)	105.9(2)	C(51)-Si(5)-C(53)	111.4(7)
P(1)-Si(1)-C(11)	109.9(5)	C(51)-Si(5)-C(54)	111.6(7)
P(1)-Si(1)-C(12)	108.7(5)	C(52)-Si(5)-C(53)	108.2(8)
P(1)-Si(1)-C(13)	114.2(5)	C(52)-Si(5)-C(54)	109.5(8)

Table 2, Continued.

C(11)-Si(1)-C(12)	109.5(8)	C(53)-Si(5)-C(54)	107.6(8)
C(11)-Si(1)-C(13)	107.3(8)	C(61)-Si(6)-C(62)	109.7(8)
C(12)-Si(1)-C(13)	107.1(8)	C(61)-Si(6)-C(63)	110.8(7)
P(1)-Si(2)-C(21)	108.1(5)	C(61)-Si(6)-C(64)	111.6(8)
P(1)-Si(2)-C(22)	112.9(6)	C(62)-Si(6)-C(63)	107.8(7)
P(1)-Si(2)-C(23)	107.9(5)	C(62)-Si(6)-C(64)	109.2(9)
C(21)-Si(2)-C(22)	109.5(8)	C(63)-Si(6)-C(64)	107.7(7)
C(21)-Si(2)-C(23)	109.0(8)	Tl(1)-C(41)-Si(4)	114.5(7)
C(22)-Si(2)-C(23)	109.3(9)	Tl(1)-C(51)-Si(5)	117.2(7)
P(1)-Si(3)-C(31)	108.1(5)	Tl(1)-C(61)-Si(6)	116.2(7)
P(1)-Si(3)-C(32)	113.5(5)		

Table 3. Non-Hydrogen Atom Fractional Coordinates and Equivalent Isotropic Thermal Parameters for **1**, with Estimated Standard Deviations in Parentheses

Atom	x	y	z	B _{iso} (Å ²) ^a
Tl	0.33779(7)	0.01648(3)	0.7786(-)	2.21(4)
P(1)	0.32523(23)	-0.01183(24)	1.0169(3)	2.11(17)
Si(1)	0.1976(3)	-0.1566(3)	1.0616(3)	2.76(20)
Si(2)	0.4544(3)	-0.0102(3)	1.0908(4)	2.97(23)
Si(3)	0.3082(3)	0.1011(3)	1.1091(3)	2.74(20)
Si(4)	0.2590(3)	-0.1886(3)	0.6235(4)	3.16(23)
Si(5)	0.1911(3)	0.1084(3)	0.6955(4)	2.85(20)
Si(6)	0.5662(3)	0.1635(3)	0.6747(4)	3.15(21)
C11	0.0902(10)	-0.1757(11)	0.9849(14)	3.8(8)
C12	0.2257(11)	-0.2520(10)	1.0249(14)	3.9(8)
C13	0.1666(12)	-0.1708(11)	1.2089(14)	4.3(9)
C21	0.4859(12)	-0.0840(13)	1.0033(13)	3.9(11)
C22	0.4335(13)	-0.0568(14)	1.2342(14)	5.0(12)
C23	0.5543(11)	0.1144(12)	1.0896(17)	5.0(11)
C31	0.1836(11)	0.0777(11)	1.0855(13)	3.5(8)
C32	0.3295(12)	0.1019(11)	1.2609(13)	3.7(9)
C33	0.3950(12)	0.2191(10)	1.0513(13)	3.3(9)
C41	0.3473(11)	-0.1126(11)	0.7290(12)	3.3(8)
C42	0.1394(11)	-0.2599(10)	0.6882(13)	3.7(9)
C43	0.2946(13)	-0.2726(15)	0.5602(17)	5.9(12)
C44	0.2450(12)	-0.1152(13)	0.5116(13)	4.8(10)
C51	0.1944(10)	0.0059(10)	0.7622(13)	3.4(7)
C52	0.0621(11)	0.0773(12)	0.6823(16)	4.9(10)
C53	0.2557(12)	0.2201(12)	0.7797(14)	4.0(9)
C54	0.2481(12)	0.1349(12)	0.5567(14)	4.4(10)
C61	0.4744(11)	0.1588(10)	0.7663(13)	3.6(8)

^aB_{iso} = the mean of the principal axes of the thermal ellipsoid

Table 3, Continued.

Atom	x	y	z	B _{iso} (Å ²) ^a
C62	0.6225(11)	0.0977(12)	0.7381(17)	5.1(11)
C63	0.6625(10)	0.2895(11)	0.6513(13)	3.7(8)
C64	0.5159(12)	0.1100(13)	0.5392(15)	4.8(10)

^aB_{iso} = the mean of the principal axes of the thermal ellipsoid

TECHNICAL REPORTS DISTRIBUTION LIST

ORGANOMETALLIC CHEMISTRY FOR ELECTRONIC & OPTICAL MATERIALS

Dr. Harold E. Guard
Code 1113
Chemistry Division, 331
Office of Naval Research
800 N. Quincy Street
Arlington, Va 22217-5660

Dr. Richard W. Drisko
Naval Facilities & Engineering
Service Center
Code L52
Port Hueneme, CA 93043

Defense Technical Information
Center
Building 5, Cameron Station
Alexandria, VA 22314

Dr. Eugene C. Fischer
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Dr. James S. Murday
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, DC 20375-5320

Dr. Bernard E. Douda
Crane Division
Naval Surface Warfare Center
Crane, IN 47522-5000

Dr. John Fischer, Director
Chemistry Division, C0235
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001

Dr. Peter Seligman
Naval Command, Control and
Ocean Surveillance Center
RDT&E Division
San Diego, CA 93152-5000